

QUVIVIQ

(daridorexant) ^{IV} 25mg, 50mg tablets

QuickStart Guide

“Taking once-nightly QUVIVIQ is the different approach I need for my insomnia.”

Lindsey Vonn
World Champion Skier &
QUVIVIQ 50 mg Patient



Individual results may vary.

Once-nightly QUVIVIQ is a **different kind of insomnia treatment**.
When taken **every night**, sleep improved over time*.

Here's what you need to know

✔ Works differently

Unlike some insomnia medications that sedate the brain, QUVIVIQ is thought to work **by turning down overactive wake signals**, one of the biological causes of insomnia.

✔ Taken differently

QUVIVIQ is designed to be **taken every night** to help you get consistent, better sleep*. In clinical studies, **stopping QUVIVIQ did not cause withdrawal symptoms**.

✔ Experienced differently

QUVIVIQ may take several nights to start working. In clinical studies, **people reported sleeping better after 1 week of taking QUVIVIQ every night**, and sleep improvements continued to increase over 3 to 4 weeks†.

*Improvements measured at months 1 and 3 in clinical studies.

†Results at month 1 were maintained at month 3.

If you're currently or have recently been taking another insomnia medication, talk to your doctor about what to expect when switching to QUVIVIQ.



To start saving on QUVIVIQ, click this button



What is QUVIVIQ (daridorexant)?

QUVIVIQ is a prescription medicine for adults who have trouble falling asleep or staying asleep (insomnia).

Important Safety Information

Do not take QUVIVIQ if you fall asleep often at unexpected times (narcolepsy) or if you are allergic to QUVIVIQ or any of its ingredients.

Please see next page for additional Important Safety Information. Please see full Prescribing Information and Medication Guide.

Important Safety Information

QUVIVIQ may cause serious side effects, including:

- **Decreased awareness and alertness.** The morning after you take QUVIVIQ, your ability to drive safely and think clearly may be decreased. You may also have sleepiness during the day.
- Do not take more QUVIVIQ than prescribed.
- Do not take QUVIVIQ unless you are able to stay in bed for at least 7 hours before you must be active again.
- Take QUVIVIQ at night within 30 minutes before going to bed.

QUVIVIQ is a federally controlled substance because it can be abused or lead to dependence.

Before taking QUVIVIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of depression, mental illness, or suicidal thoughts or actions; drug or alcohol abuse or addiction; a sudden onset of muscle weakness (cataplexy); daytime sleepiness
- have lung or breathing problems, including sleep apnea
- have liver problems
- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Taking QUVIVIQ with certain medicines can cause serious side effects. QUVIVIQ may affect the way other medicines work and other medicines may affect the way QUVIVIQ works.
- **Do not take QUVIVIQ with other medicines that can make you sleepy unless instructed by your healthcare provider.**

What should I avoid while taking QUVIVIQ?

- Do not drink alcohol while taking QUVIVIQ. It can increase the effects of alcohol, which can be dangerous.
- Do not drive, operate heavy machinery, do anything dangerous, or do other activities that require clear thinking if you do not feel fully awake, or you have taken QUVIVIQ and have less than a full night of sleep (at least 7 hours), or if you have taken more QUVIVIQ than prescribed.

QUVIVIQ may cause other serious side effects, including:

- **Worsening depression and suicidal thoughts.** Call your healthcare provider right away if you have any worsening depression or thoughts of suicide or dying.
- **Temporary inability to move or talk (sleep paralysis) for up to several minutes, or hallucinations while you are going to sleep or waking up.**
- **Complex sleep behaviors** such as sleep-walking, sleep-driving, preparing and eating food, making phone calls, having sex or doing other activities while not fully awake that you may not remember the next morning. Stop taking QUVIVIQ and call your healthcare provider right away if you experience a complex sleep behavior.

The most common side effects of QUVIVIQ are headache and sleepiness.

These are not the only side effects of QUVIVIQ. Call your doctor for advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information and Medication Guide.

